

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



Preliminary Result of the Occultation by (3200) Phaethon as Observed at Ballarat, California

Quanzhi Ye (叶泉志)^{1,2,3} , Qicheng Zhang⁴ , Man-To Hui (許文韜)^{5,6} , and David Dunham⁷

Published 2019 December 12 • © 2019. The American Astronomical Society. All rights reserved.

Research Notes of the AAS, Volume 3, Number 12

qye@umd.edu

¹ Division of Physics, Mathematics and Astronomy, California Institute of Technology, Pasadena, CA 91125, USA

² Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA

³ Department of Astronomy, University of Maryland, College Park, MD 20742, USA

⁴ Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125, USA

⁵ Department of Earth, Planetary and Space Sciences, UCLA, 595 Charles Young Drive East, Los Angeles, CA 90095, USA

⁶ Institute for Astronomy, University of Hawai'i, Honolulu, HI 96822, USA

⁷ IOTA, 3719 Kara Ct., Greenbelt, MD 20770, USA

Quanzhi Ye <https://orcid.org/0000-0002-4838-7676>

Qicheng Zhang <https://orcid.org/0000-0002-6702-191X>

Man-To Hui <https://orcid.org/0000-0001-9067-7477>

Received 2019 November 26

Accepted 2019 December 11

Published 2019 December 12

Quanzhi Ye *et al* 2019 *Res. Notes AAS* **3** 188

<https://doi.org/10.3847/2515-5172/ab60aa>

Asteroid occultation ; Near-Earth objects ; Occultation

Export citation and abstract

BibTeX

RIS

Near-Earth asteroid (3200) Phaethon is the primary target for the Demonstration and Experiment of Space Technology for INterplanetary voYage, Phaethon fLyby and dUst Science Phaethon fLyby with reUSable probe (DESTINY+) mission, currently being developed by the Japan Aerospace Exploration Agency (JAXA). The size of Phaethon is measured to be about 5–6 km, although numbers derived by different techniques are not in strict agreement: radar measurement suggests a sphere with a diameter of 6.2 km, or a top-shaped figure with equivalent spherical diameter of 5.5 km (Taylor et al. 2019), while two thermophysical models based on infrared data suggest a spherical diameter of 5.1 ± 0.2 km (Hanuš et al. 2016) and $4.6^{+0.2}_{-0.3}$ km (Masiero et al. 2019), $\gtrsim 2\sigma$ different from radar data.

On UT 2019 July 29, Phaethon passed in front of a 7.3 mag star SAO 40261, providing an unique opportunity to constrain Phaethon's size using a technique other than radar or thermal observation. The occultation was visible in a narrow strip from southern Colorado to central California. Guided by the prediction made by Southwest Research Institute (SwRI) based on JPL orbit solution #701,⁸ we traveled to the deserted town of Ballarat, California to observe this occultation. According to the prediction, the center line would pass about 1.5 km north of the town center. We set up two telescopes, an Astro-Tech 72 mm refractor and an Astro-Tech 60 mm refractor, at $36^{\circ}2'58''.9$ N, $117^{\circ}13'34''.3$ W, ~ 1 km south of the predicted center line, though preliminary post-occultation analysis showed that this site is within ~ 100 m of the true center line (S. Preston 2019, private communication). The telescopes were about 25 m apart, with the 72 mm telescope to the south and the 60 mm telescope to the north, in a direction that was roughly perpendicular to the motion of Phaethon's shadow. For each telescope, we used a QHY CMOS camera to record the event at a frame rate of ~ 30 frames per second.

The "south" telescope successfully recorded the occultation; the "north" telescope did not collect any useful data due to a pointing issue. Figure 1 shows the light-curve derived from the data collected by the "south" telescope. The duration of the occultation is 0.498 ± 0.004 s. Considering the speed of Phaethon's shadow to be 11.15 km s^{-1} , corrected for Earth's rotation at Ballarat, the length of the shadow is 5.55 ± 0.05 km, in agreement with the radar

measurement. We note that this does not automatically reject the thermophysical models, as we could be measuring the long axis of Phaethon. A detailed analysis of this occultation event, using the full set of observations collected by International Occultation Timing Association members and a SwRI team, is underway.




Figure 1. Light-curve of occultation as recorded by the "south" telescope. Times are in UT and have an absolute error of 0.03 s. A time-lapse video of the occultation event is available online.

We thank SwRI for the prediction, as well as H. Pavlov for helpful discussion on the basics of occultation observation. This research used equipment of the Cahill Rooftop Observatory, managed and funded by Department of Astronomy, California Institute of Technology, and is partially supported by a NASA grant.

Footnotes

8 <http://iota.jhuapl.edu/20190729PhaethonBefore.htm>

References


↑ Hanuš J., Delbo' M., Vokrouhlický D. *et al* 2016 *A&A* **592** A34
Crossref ADS Google Scholar



↑ Masiero J. R., Wright E. L. and Mainzer A. K. 2019 *AJ* **158** 97
IOPscience (<https://iopscience.iop.org/1538-3881/158/3/97>)
Scholar



ADS Google

 Taylor P. A., Rivera-Valentín E. G., Benner L. A. M. *et al* 2019 *P&SS* **167** 1

[Crossref](#)

[ADS](#)

[Google Scholar](#)



Export references:

[BibTeX](#)

[RIS](#)